




MAYO SCORING INDEX – ASSESSMENT FOR ULCERATIVE COLITIS ACTIVITY


Mayo Score Disease Activity Index (DAI)	Answer choices	Pts
Stool Frequency	Normal number of stools	0
	1–2 stools/day more than normal	1
	3–4 stools/day more than normal	2
	>4 stools/day more than normal	3
Rectal bleeding *	None	0
	Visible blood with stool less than half the time	1
	Visible blood with stool half of the time or more	2
	Passing blood alone	3
Endoscopic Findings **	Normal or inactive disease	0
	Mild disease (erythema, decreased vascular pattern, mild friability)	1
	Moderate disease (marked erythema, absent vascular pattern, friability erosions)	2
	Severe disease (spontaneous bleeding, ulceration)	3
Physician Global Assessment	Normal	0
	Mild colitis	1
	Moderate colitis	2
	Severe colitis	3

ENDOSCOPIC FINDINGS

0


1


2


3


* A score of 3 for bleeding required patients to have at least 50 % of bowel motions accompanied by visible blood and at least one bowel motion with blood alone

** The mucosal appearance at endoscopy is not included in the Partial Mayo Score

Ref.: Schroeder KW, Tremaine WJ, Ilstrup DM. N Engl J Med. 1987; 317(26):1625-9